

ORIGINAL ARTICLE

**IMPAIRED ADULT PERFORMANCE AND REPRODUCTION OF THE PINK BOLLWORM
Pectinophora gossypiella (SAUNDERS)(LEPIDOPTERA: GELECHIDAE) AFTER TREATMENT
OF EGGS WITH THE CHITIN SYNTHESIS INHIBITORS, NOVIFLUMURON AND
NOVALURON.**

M. Tanani and *K. Ghoneim

Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt.

Article History: Received 5th February ,2017, Accepted 1st March,2017, Published 2nd March,2017

ABSTRACT

The present study was conducted aiming to investigate the affected adult performance and reproductive potential of the pink bollworm *P. gossypiella*, a dangerous pest of cotton worldwide, after treatment of 1-day old eggs with sublethal concentrations of Noviflumuron (4.0, 2.0, 1.0 and 0.5 ppm) and Novaluron (1.0, 0.5, 0.1 and 0.05 ppm). The adult emergence was considerably blocked by both compounds but they failed to affect the adult survival or morphogenesis. The total adult longevity was remarkably shortened. Noviflumuron treatment caused a slight prolongation or shortening of pre-oviposition period, depending on the concentration but Novaluron treatment led to slightly prolonged period. The oviposition period was remarkably shortened. Pronouncedly prolonged post-oviposition period was recorded by Noviflumuron but a slightly prolonged period by Novaluron. The oviposition efficiency was conspicuously prohibited, regardless the tested compound. Fecundity of the successfully reproducing females and fertility of eggs laid by these females were dramatically reduced. The incubation period of eggs laid by the females was considerably prolonged, regardless the tested compound.

Keywords: Emergence, Fecundity, Fertility, Incubation, Longevity, Oviposition.

1.INTRODUCTION

The pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae), is one of the most destructive insects attacking cotton fields world-wide, and a particularly pest difficult to control (Liu *et al.*, 2009). It has known to cause serious loss in seed cotton yield, oil content and normal opening of bolls (Patil, 2003) as well as the grade and staple of the lint (Mohammed, 2013). Further the cotton lint in the insect infested bolls gets damaged by secondary fungal infection (Kranthi, 2015). This insect pest attacks cotton plants in Egypt during the flowering and fruiting stages (Khidr *et al.*, 1996) causing enormous reduction in quantity and quality of cotton yield arising to one million kantar annually when it neglected (El-Aswad and Aly, 2007; Khatter and Abuldahb, 2011; Kandil *et al.*, 2012). In Egypt, control of *P. gossypiella* depends mostly on the use of various conventional insecticides (Radwan and El-Malla, 2015). It has recently developed high levels of resistance against most of those synthetic compounds currently used in cotton fields because of its ability to detoxify these chemicals (Abd-Elhady

and Abd El-Aal, 2011; Sabry and Abdel-Aziz, 2013). Moreover, it has been reported to develop resistance against the transgenic cotton varieties in Arizona (USA)(Fabrick and Tabashnik, 2012) and different districts of India (Monsanto, 2010). In addition to the development of resistance, the intensive and discriminate uses of many conventionally synthetic pesticides led to several drastic problems, such as the environmental pollution, hazards to human and animals like birds, fishes and mammals, destruction of the natural enemies, like parasites, predators (Rose, 2001; Rashad *et al.*, 2015). Therefore, alternative materials have been initiated recently to minimize the pesticide hazards (Derbalah *et al.*, 2014) and to delay the resistance development in *P. gossypiella* (Salama *et al.*, 2013).

During the late few decades, a new class of comparatively safe compounds have been developed and known as insect growth regulators (IGRs) (Dhadialla *et al.*, 1998; Khan and Qamar, 2012). In contrast to the classical chemical insecticides, IGRs are not directly toxic, but act selectively on the development, metamorphosis or reproduction of the target insect species (Hoffmann and Lorenz, 1998; Martins and Silva, 2004). They are quite selective in their mode of action and potentially act only on the target species (Sabry and Abdou, 2016). Chitin synthesis inhibitors (CSIs) are usually classified in IGRs (Tunaz and Uygun, 2004) interfering with

*Corresponding author: **Dr. K. Ghoneim**, Department of Zoology and Entomology, Faculty of Science, Al-Azhar University, Cairo, Egypt

chitin biosynthesis in insects and thus prevents moulting, or produces an imperfect cuticle (Hammock and Quistad, 1981). These compounds affect, also, the hormonal balance in insects, thereby resulting in physiological disturbances (Soltani *et al.*, 1984).

Noviflumuron is a new chemistry currently being developed by Dow AgroSciences, Indianapolis, Indiana, USA, for the structural pest control market (Ameen *et al.*, 2002). Noviflumuron is a benzoylphenyl urea CSI that prevents the successful molting and development of some insects, such as fleas, ants, termites and houseflies (Sheets and Karr, 2001; Karr *et al.*, 2004). Its suspension concentrate, dust or gel bait can effectively suppress *Blattella germanica* populations with a pattern of activity similar to that expected from a CSI (Ameen *et al.*, 2005; Smith *et al.*, 2002; Wang and Bennett, 2006). Also, adult *B. germanica* exposed to Noviflumuron fail to produce viable eggs, hence their reproductive potential is reduced (King, 2005). Novaluron is a relatively new benzoylphenyl urea CSI with low mammalian toxicity (Barazani, 2001; Ishaaya and Horowitz, 2002). The compound has no appreciable effect on parasitoids and has probably a mild effect on the natural enemies (Ishaaya *et al.*, 2001, 2002). Its residues tend to dissipate with half-life of 2.08 days and the safe use of it on tomatoes, and possibly on other crops in Egypt was established (Malhat *et al.*, 2014). Novaluron is a powerful suppressor of the pest populations, such as *Bemisia tabaci* and *Trialeurodes vaporariorum* (Ishaaya *et al.*, 2003). It acts by ingestion and contact against several insect pests, such as *Spodoptera* spp., *Tuta absoluta*, *Helicoverpa armigera*, and *Liriomyza huidobrensis* (Kim *et al.*, 2000). It exhibited, also, a good activity against the Colorado potato beetle (Cutler *et al.*, 2005 a,b, 2007; Alyokhin *et al.*, 2009) and impaired the development (Ghoneim *et al.*, 2015) and adult performance (Hamadah *et al.*, 2015) of *Spodoptera littoralis*. Recently, treatment of newly hatched and full grown larvae of *P. gossypiella* with Novaluron led to reduced survival, retarded development, impaired metamorphosis (Ghoneim *et al.*, 2017a), disrupted adult performance, inhibited reproductive potential (Hassan *et al.*, 2017), declined main metabolites (Tanani *et al.*, 2017), and deteriorated larval haemogram (Ghoneim *et al.*, 2017b). According to the available literature, few studies had been conducted investigating the ovicidal effects of insecticides and novel IGRs on eggs of lepidopterous pests but the majority focused on treatment of larval stage (Abd-El-Aziz and Sayed, 2014). Taking all of these considerations into account, the present study was carried out to evaluate the long-term effects of Noviflumuron and Novaluron, novel CSIs, on the adult performance and reproductive potential of *P. gossypiella* after treatment of 1-day old eggs.

2. MATERIALS AND METHODS

Experimental Insect

A culture of the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) was originated by a sample of newly hatched larvae from the susceptible culture maintained for several generations along some years in Plant Protection Research Institute, Doqqi, Giza, Egypt. It was reared under constant conditions (27±2°C and 75±5% R.H.) at Department of Zoology and Entomology, Faculty of science, Al-Azhar University, Cairo. Larvae were provided

with an artificial diet as described by Abd El-Hafez *et al.* (1982). For rearing details and manipulation of all developmental stages under the laboratory controlled conditions, see Ghoneim *et al.* (2017a).

CSIs and Preparation of Concentrations

The tested Benzoylphenylurea compounds, in the present study, are Noviflumuron and Novaluron. Noviflumuron (Recruit III, and Recruit III AG) has the chemical name: N{[[[3,5-dichloro-2-fluoro-4-(1,1,2,3,3,3 hexafluoropropoxy) phenyl] amino]carbonyl]-2,6-difluorobenzamide. Its molecular formula is C₁₇H₇C₁₂F₉N₂O₃. It was kindly obtained from Laboratory of Pesticides, Agricultural Research Center, Doqqi, Giza, Egypt. Novaluron (Rimon) has the chemical name: [1-[chloro-4-(1,1,2-trifluoromethoxyethoxy) phenyl] -3-(2,6-difluorobenzoyl) urea]. Its molecular formula is C₁₇H₉C₁F₈N₂O₄. It was purchased from Sigma-Aldrich Chemicals. Four concentrations of each CSI were prepared by diluting with distilled water in volumetric flasks as follows: Noviflumuron: 4.0, 2.0, 1.0 and 0.5 ppm. Novaluron: 1.0, 0.5, 0.1 and 0.05 ppm.

Egg Treatment

Four groups of freshly emerged moths of *P. gossypiella*, each group 10 pairs (♂ X ♀), were confined in a glass chimney cage (17 cm height and 7.12 cm in diameter), inside which a piece of cotton wool previously soaked in 20% sugar solution was suspended to be renewed 48 hr for moths' nutrition (El-Barkey *et al.* 2009). After deposition of eggs on plastic papers at the top and bottom of the cage, these papers had been divided into four pieces in order to obtain four replicates of 1-day old eggs (10 eggs/replicate). Then, each replicate was carefully sprayed with each concentration of each CSI using an atomizer. All egg papers were left for 15 min. to allow evaporation of excess water under laboratory conditions. Similar replicates of control eggs were treated with distilled water only using the same technique.

Each of the CSI-treated and control replicates was kept separately in a suitable plastic jar for hatching under the controlled conditions (27±2°C and 75±5% R.H.). After the incubation period, each newly hatched larva was transferred into a glass tube (6.0x1.5 cm) containing 2 g of artificial diet. The treated and control larvae and pupae were carefully handled until the adult emergence. Newly emerged moths were sexed and transferred to chimney glass cage (six pairs /cage). The moths were fed on 20% sucrose solution. Cages were examined daily to record all parameters of adult performance and reproductive criteria.

Adult Performance Parameters

Adult emergence: Number of emerged adults was expressed in % according to Jimenez-Peydro *et al.* (1995) as follows:

$$[\text{No. of completely emerged adults} / \text{No. of pupae}] \times 100$$

Adulticidal activity: Adulticidal activity of the tested CSIs was detected by mortalities throughout the adult longevity and calculated in percentage.

Morphogenic efficiency: The morphogenic efficiency of the tested CSIs was detected by the adult deformities and calculated in percentage as follows:

$$[\text{No. of deformed adults} / \text{No. of emerged adults}] \times 100$$

Adult longevity: Total adult longevity of females was measured in mean days±SD. The major compartments of

adult longevity are pre-oviposition (ovarian maturation) period, oviposition period (reproductive life-time) and post-oviposition period. All durations were measured in mean days \pm SD.

Reproductive Criteria

Oviposition rate was calculated as follows:

Number of laid eggs per ♀/reproductive lifetime (in days). Eggs laid by the successfully reproducing adult females were counted for calculating the number of eggs per female (Fecundity). The laid eggs were kept in Petri dishes under the same controlled laboratory conditions as previously mentioned. Just after the oviposition, eggs were observed until hatching elapsing an incubation period (in days). The hatchability (Fertility) was usually expressed in hatching percentage of laid eggs. Sterility index was calculated according to Topozada *et al.* (1966) as follows:

$$\text{Sterility Index} = 100 - [(a \ b / A \ B) \times 100]$$

Where: a: mean number of eggs laid per female in the treatment. b: percentage of hatching in the treatment. A: mean number of eggs laid per female in the controls. B: percentage of hatching in the controls.

Statistical analysis of data

Data obtained were analyzed by the Student's *t*-distribution, and refined by Bessel correction (Moroney, 1956) for the test significance of difference between means.

3. RESULTS

Effects of CSIs on Adult Performance of *P. gossypiella*

After treatment of 1-day old eggs of *P. gossypiella* with four sublethal concentrations (4.0, 2.0, 1.0 and 0.5 ppm) of Noviflumuron, data of the most informative parameters of adult performance were arranged in Table (1). Depending on these data, the adult emergence was considerably blocked, in a dose-dependent course (90.0, 88.25, 88.17 and 78.44 emergence %, at 0.5, 1.0, 2.0 and 4.0 ppm, respectively, vs. 100% emergence of control adult females). Noviflumuron failed to exhibit neither adulticidal nor morphogenic effect, since no mortality or adult deformity had been observed. With regard to the adult longevity and its main compartments, data contained in the same table clearly revealed that the total longevity of adult females was remarkably shortened, in a dose-dependent manner (13.7 \pm 0.96, 13.5 \pm 0.71, 13.2 \pm 0.50 and 10.5 \pm 0.80 days, at 0.5, 1.0, 2.0 and 4.0 ppm, respectively, in comparison with 15.0 \pm 0.82 days of control adult females). In respect of the pre-oviposition period (ovarian maturation period), treatment of 1-day old eggs with the highest concentration level led to failure of the successfully emerged adult females to lay eggs, thus the pre-oviposition period could not be measured. However, such period was affected after treatment of eggs with other concentrations, in no certain trend (3.5 \pm 1.29, 1.5 \pm 0.71 and 2.5 \pm 1.80 days, at 0.5, 1.0 and 2.0 ppm, respectively, vs. 2.3 \pm 0.50 days of control females). Moreover, Noviflumuron exerted a tremendously enforcing action on the ovipositing adult females to quickly lay eggs, since their oviposition period (reproductive life-time) was significantly shortened, in a reverse course of concentration (8.7 \pm 0.90, 8.5 \pm 0.71 and 8.0 \pm 1.41 days, at 2.0, 1.0 and 0.5 ppm, respectively, compared to 11.3 \pm 0.96 days of control females). The last compartment of adult longevity is post-oviposition period which was pronouncedly prolonged, in no certain trend (3.3 \pm 0.77, 3.5 \pm 0.71 and 2.2 \pm 0.50 days, at

2.0, 1.0 and 0.5 ppm, respectively, vs. 1.5 \pm 1.29 days of control females).

After treatment of 1-day old eggs of *P. gossypiella* with four sublethal concentrations (1.0, 0.5, 0.1 and 0.05 ppm) of Novaluron, data of the adult life parameters were assorted in Table (2). On the basis of these data, the adult emergence was remarkably blocked, in a dose-dependent course (93.80, 87.50, 75.88 and 75.50 emergence %, at 0.05, 0.1, 0.5 and 1.0 ppm, respectively, vs. 100% emergence of control adult females). Novaluron failed to affect the survival of successfully emerged adult females or their morphogenesis, since no mortality or adult malformations had been observed. In connection with the adult longevity and its main compartments, data included in the same table exiguously demonstrated that the total longevity was elaborately shortened only at the highest concentration level (11.8 \pm 0.50 days, compared to 14.8 \pm 0.50 days of controls) but slightly shortened at other concentrations, in no certain trend. After treatment of eggs with the highest concentration level of Novaluron, the emerged adult females failed to lay eggs, thus no pre-oviposition period could be estimated. At other concentrations, the pre-oviposition period was unremarkably prolonged, in a dose-dependent fashion (2.5 \pm 0.58, 2.8 \pm 0.58 and 2.9 \pm 0.96 days, at 0.05, 0.1 and 0.5 ppm, respectively, vs. 2.3 \pm 0.50 days of control adult females), indicating a slight retarding effect of Novaluron on the ovarian maturation rate. On the other hand, the oviposition period was seriously shortened, in a dose-dependent course (10.3 \pm 1.71, 9.0 \pm 0.82 and 8.5 \pm 1.29 days, at 0.05, 0.1 and 0.5 ppm, respectively, vs. 11.0 \pm 0.82 days of control congeners) denoting a powerful inducing action of Novaluron on the ovipositing females to lay eggs quickly. Also, data of the aforementioned table clearly displayed that the post-oviposition period was slightly prolonged, in a dose-dependent trend 2.0 \pm 0.82, 2.3 \pm 0.96 and 3.0 \pm 1.41 days, at 0.05, 0.1 and 0.5 ppm, respectively, vs. 1.5 \pm 0.58 days of control congeners).

For the comparative purpose, there is no remarkable difference in effect on Noviflumuron and Novaluron on adult survival, morphogenesis or blocking emergence. However, Noviflumuron exhibited a relatively stronger shortening effect than Novaluron on the total adult longevity and oviposition period.

Effects of CSIs on the Reproductive Potential of *P. gossypiella*

After treatment of 1-day old eggs of *P. gossypiella* with sublethal concentrations of Noviflumuron, data of the most important reproductive criteria had been distributed in Table (3). Depending on these data, the highest concentration level could be described as the extremely anti-reproductive one because the successfully emerged adult females failed to lay eggs in spite of the mating with normal adult males. The oviposition efficiency of treated females was tremendously prohibited by Noviflumuron, since the oviposition rate was drastically regressed, in a dose-dependent course (3.4 \pm 1.50, 3.1 \pm 0.49 and 2.2 \pm 1.20, at 0.5, 1.0 and 2.0 ppm, respectively, vs. 15.4 \pm 1.45 of control congeners). Data of the same table evidently revealed that the fecundity (mean number of eggs/♀) was dramatically reduced, in a consecutive correlation with the concentration (28 \pm 14.90, 26 \pm 2.12 and 20 \pm 2.30 eggs/treated ♀, at 0.5, 1.0 and 2.0 ppm, respectively, vs. 174 \pm 24.39 eggs/control ♀). Another parameter of the

reproductive capacity is fertility (hatching % of laid eggs or egg viability) which was severely reduced, in a dose-dependent manner (38.4, 29.4 and 10.0%, at 0.5, 1.0 and 2.0 ppm, respectively, vs. 71.2% of control hatchability). In other words, the sterility index considerably increased as the concentration was increased. With regard to the embryonic development in the laid eggs, it was seriously retarded by Noviflumuron, as detected by the remarkably prolonged incubation period (4.8±0.50, 5.5±0.71 and 5.8±0.50 days, at 0.5, 1.0 and 2.0 ppm, respectively, vs. 4.3±0.0 days of laid eggs by control females).

After treatment of 1-day old eggs of *P. gossypiella* with the sublethal concentrations of Novaluron, data of the most important reproductive criteria were summarized in Table (4). As obviously shown in this table, no oviposition was observed for the adult females after treatment of 1-day old eggs with the highest concentration. Thus, no reproductive data could be recorded. At other concentrations, the oviposition efficiency was tremendously inhibited by Novaluron, since the oviposition rate was severely regressed, in a dose-dependent course (10.2±5.02, 7.3±1.55 and

4.8±1.18, at 0.05, 0.1 and 0.5 ppm, respectively, vs. 19.7±4.27 of control congeners). Also, fecundity was drastically reduced, proportional to the concentration (109±57.19, 65±11.34 and 41±12.19 eggs/treated ♀, at 0.05, 0.1 and 0.5 ppm, respectively, vs. 218±56.12 eggs/control ♀). Sterility index considerably increased as the concentration was increased. In addition, data of the same table clearly displayed that Novaluron exerted a strong retarding action on the embryonic development, since the incubation period was significantly prolonged, in a dose-dependent course (4.8±0.5, 4.8±0.50 and 4.9±0.58 days, at 0.05, 0.1 and 0.5 ppm, respectively, vs. 4.3±0.5 days of eggs laid by control females).

For the comparative purpose, the deteriorating effect of Noviflumuron on the oviposition rate, fecundity and fertility appeared to be stronger than that exhibited by Novaluron. In respect of the incubation period, and subsequently the embryonic development, there was no considerable difference between the potencies of the tested CSIs.

Table 1 adult performance of *P. gossypiella* as affected by treatment of 1- day old eggs with Noviflumuron

Conc. (ppm)	Adult emergence (%)	Adult mortality (%)	Adult deformations (%)	Adult longevity (mean days ±SD)			Total longevity
				Ovarian maturation period	Reproductive lifetime	Post-oviposition period	
4.0	78.44	00.0	00.0	---	---	---	10.5±0.80 c
2.0	88.17	00.0	00.0	2.5±1.80 a	8.7±0.90 b	3.3±0.77 b	13.2±0.50 b
1.0	88.25	00.0	00.0	1.5±0.71 a	8.5±0.71 b	3.5±0.71 b	13.5±0.71 b
0.5	90.0	00.0	00.0	3.5±1.29 a	8.0±1.41 c	2.2±0.50 b	13.7±0.96 a
Control	100.0	00.0	00.0	2.3±0.50	11.3±0.96	1.5±1.29	15.0±0.82

Conc.: concentration level. Mean±SD followed with a: not significantly different (P>0.05), b: significantly different (P<0.05), c: highly significantly different (P<0.01). --- : adult females failed to lay eggs although mating with normal males was done.

Table 1 adult performance of *P. gossypiella* as affected by treatment of 1- day old eggs with Novaluron

onc. (ppm)	Adult emergence (%)	Adult mortality (%)	Adult deformations (%)	Adult longevity (mean days ±SD)			Total longevity
				Ovarian maturation period	Reproductive life-time	Post-oviposition period	
1.0	75.50	00.0	00.0	---	---	---	11.8±0.50 b
0.5	75.88	00.0	00.0	2.9±0.96 a	8.5±1.29 b	3.0±1.41 a	14.3±0.96 a
0.1	87.50	00.0	00.0	2.8±0.58 a	9.0±0.82 b	2.3±0.96 a	13.8±0.96 a
0.05	93.80	00.0	00.0	2.5±0.58 a	10.3±1.71 b	2.0±0.82 a	14.8±0.50 a
Control	100.0	00.0	00.0	2.3±0.50	11.0±0.82	1.5±0.58	14.8±0.50

Conc., a, b, ---: see footnote of Table 1.

Table 3. Reproductive potential of *P. gossypiella* adults as affected by treatments of 1-day old eggs with Noviflumuron.

Conc. (ppm)	Oviposition rate	Fecundity (No. of eggs/♀±SD)	Fertility (%)	Sterility index (%)	Incubation period (mean days±SD)
4.0	0.0	---	---	---	---
2.0	2.2±1.20 d	20±2.30 c	10.0	98.38	5.8±0.50 b
1.0	3.1±0.49 d	26±2.12 c	29.4	93.83	5.5±0.71 b
0.5	3.4±1.50 d	28±14.90 d	38.4	91.32	4.8±0.50 b
Control	15.4±1.45	174±24.39	71.2	---	4.3±0.50

Conc., a, b, c, ---: see footnote of Table 1. d: very highly significantly different (P<0.001).

Table 4. Reproductive potential of *P. gossypiella* adults as affected by treatments of 1-day old eggs with Novaluron.

Conc. (ppm)	Oviposition rate	Fecundity (mean No. of eggs/♀±SD)	Fertility (%)	Sterility index (%)	Incubation period (mean days ±SD)
1.0	0.0	---	---	---	---
0.5	4.8±1.18 d	41±12.19 d	31.1	91.89	4.9±0.58 b
0.1	7.3±1.55 c	65±11.34 c	45.3	81.27	4.8±0.50 b
0.05	10.2±5.02 b	109±57.19 b	49.8	65.46	4.8±0.50 b
Control	19.7±4.27	218±56.12	72.1	---	4.3±0.50

Conc., a, b, c, ---: see footnote of Table 1. d: see footnote of Table 3.

4. DISCUSSION

Influenced Adult Performance of *P. gossypiella* by CSIs

Blocked adult emergence

According to the currently available literature, the majority of studies focused on the investigation of the effects of IGRs after treatment of larvae. The adult emergence of different insect species was significantly hindered after treatment of larvae with several IGRs (including CSIs), such as *Plutella xylostella* by Hexaflumuron (Mahmoudvand *et al.*, 2012); *Drosophila melanogaster* by Pyriproxyfen (Benseba *et al.*, 2015); *Spodoptera littoralis* by Novaluron (Ghoneim *et al.*, 2015) or Cyromazine (Tanani *et al.*, 2015); *Glyphodes pyloalis* by Lufenuron (Aliabadi *et al.*, 2016); *Culex quinquefasciatus* and *Aedes albopictus* by Pyriproxyfen and Methoprene (Khan *et al.*, 2016); *P. gossypiella* by Novaluron (Hassan *et al.*, 2017). Moreover, adult emergence was completely blocked in *Corcyra cephalonica* after treatment of 4th instar larvae with Fenoxycarb (Singh and Tiwari, 2016). In addition, treatment of *Encarsia formosa* pupae with Pyriproxyfen resulted in prohibited adult emergence (Wang and Liu, 2016).

On the other hand, very scarce results of blocked adult emergence had been reported after treatment of eggs, such as the blocked adult emergence of *P. gossypiella* after treatment of pre-hatching eggs with LC₅₀ of Hexaflumuron (El-Barkey *et al.*, 2009). In agreement with those reported results, adult emergence of *P. gossypiella*, in the present study, was drastically blocked after treatment of 1-day old eggs with sublethal concentrations of Noviflumuron and Novaluron, in a dose-dependent course. The present result of blocked adult emergence can be interpreted by the interference of the tested CSIs with some aspects of the hormonal regulation, such as disturbance of release of adult eclosion hormone and/or

inhibition of the neurosecretion (prothoracicotropic hormone, PTH)(Al-Sharook *et al.*, 1991; Josephraj Kumar *et al.*, 1999).

Affected adult survival

Depending on the available literature, adulticidal effects of IGRs on some insect species are very scarcely reported, such as *S. littoralis* after treatment of larvae with the Novaluron, especially at the higher concentrations (Hamadah *et al.*, 2015), *Delia antique* after treatment of larvae with Pyriproxyfen (Zhou *et al.*, 2016) and *P. gossypiella* after treatment of the newly hatched larvae with Novaluron (Hassan *et al.*, 2017). Results of the present study were inconsistent with those reported results, since neither Noviflumuron nor Novaluron could exhibit an adulticidal effect on *P. gossypiella*, after treatment of 1-day old eggs. This failure of toxicity on adults can be understood in the light of the following suggestions. Noviflumuron and Novaluron may be metabolized or chemically degraded throughout the developmental stages of *P. gossypiella*, and subsequently no retention or distribution could be taken place in the insect body. Also, these CSIs may be denied or challenged by a high detoxification capacity of adults.

Impaired adult morphogenesis

Impaired adult morphogenesis, as expressed in the production of deformed adults, was widely reported in the available literature after treatment of larvae of various insects with different IGRs (or CSIs), such as *S. littoralis* by Tebufenozide and methoxyfenozide (Pineda *et al.*, 2004), Flufenoxuron (Bakr *et al.*, 2010), Novaluron (Hamadah *et al.*, 2015); *Rhynchophorus ferrugineus* by Diofenolan (Tanani, 2001); *Eurygaster integriceps* by Pyriproxyfen (Mojaver and Bandani, 2010); *Dysdercus koenigii* by Flucycloxuron (Khan

and Qamar, 2011); *Anagasta kuehniella* by Diflubenzuron and hexaflumuron (Ashouri *et al.*, 2014); *Helicoverpa armigera* with Hexaflumuron (Taleh *et al.*, 2015); *C. cephalonica* by Fenoxycarb (Begum and Qamar, 2016); *etc.*

Results of the present study disagreed with the previously reported results, since both Noviflumuron and Novaluron failed to affect the adult morphogenesis of *P. gossypiella*, after treatment of 1-day old eggs with sublethal concentrations. Also, the present result contradictory to those reported results of the production of deformed adults of the same lepidopterous insect, after treatment of eggs of different ages with Hexaflumuron (El-Barkey *et al.*, 2009) or after treatment of 1-day old eggs with LC₅₀ values of Lufenuron, Chlorfluazuron and Chromafenozide (Kandil *et al.*, 2012). On the contrary, the present result coincided with the failure of Novaluron to affect the adult morphogenesis of the same lepidopterous insect (Hassan *et al.*, 2017). However, the disappearance of adult deformities of *P. gossypiella*, in the current investigation, may be explained by the failure of Noviflumuron and Novaluron to disturb the hormonal regulation of the adult metamorphosis program.

Disturbed adult longevity

Total adult longevity: After the attainment of sexual maturity, insects often show degenerative changes in some tissues and organs which can be called 'senility' or 'aging'. In insects, the affected adult longevity can be considered an informative indicator for the adult aging, i.e., prolongation of longevity may denote a delay of aging and *vice versa*. In the present study, the total adult longevity of *P. gossypiella* was remarkably shortened after treatment of 1-day old eggs with sublethal concentrations of Noviflumuron or Novaluron. Comparatively, Noviflumuron exhibited a stronger shortening effect than Novaluron. The present result was in accordance with those reported results for some insects by different IGRs, such as *S. littoralis* by Lufenuron (Sammour *et al.*, 2008), Methoxyfenozide (Pineda *et al.*, 2009) and Novaluron (Hamadah *et al.*, 2015); *Agrotis ipsilon* by Flufenoxuron (El-Sheikh, 2002); *Grapholita molesta* (Reinke and Barrett, 2007) and *Spodoptera exigua* (Luna *et al.*, 2011) by Methoxyfenozide and *G. pyloalis* by Lufenuron (Aliabadi *et al.*, 2016). Also, the present result of considerably shortened total adult longevity of *P. gossypiella*, as a response to treatment of 1-day old eggs with Noviflumuron and Novaluron, corroborated with several reported results for the same lepidopterous insect after treatment of newly hatched larvae with Diflubenzuron (Kandil *et al.*, 2005; Rashad *et al.*, 2006; Salem, 2015), Chlorfluazuron (Kandil *et al.*, 2005), Chromafenozide (Salem, 2015) and Methoxyfenozide (Sabry and Abdou, 2016).

On the contrary, the present result was inconsistent with those reported results of prolonged adult longevity of the same lepidopterous insect after treatment of eggs of different ages with Hexaflumuron (El-Barkey *et al.*, 2009), after treatment of 1-day old eggs with LC₅₀ values of Lufenuron, Chlorfluazuron and Chromafenozide (Kandil *et al.*, 2012), as well as after treatment of newly hatched larvae with Hexaflumuron and Chlorfluazuron (Kandil *et al.*, 2013), Lufenuron and Pyriproxyfen (Sabry and Abdou, 2016). However, no effect was exhibited by some IGRs on the total adult longevity of some insects, such as Diufenolan against

Musca domestica (Hamadah, 2003), Tebufenozide or methoxyfenozide against *Cydia pomonella* (Saenz-de-Cabezón *et al.*, 2005), Buprofezin against *S. littoralis* (Ragaei and Sabry, 2011) and Novaluron against *Lygus lineolaris* (Portilla *et al.*, 2012).

To explain the predominantly shortened adult longevity of *P. gossypiella*, in the current study, the tested CSIs might exert a general accelerating action on these adult females to quickly pass aging ending in death. Noviflumuron exerted a stronger accelerating action than Novaluron. However, this result can be interpreted by the accumulation of xenobiotics in the body which upsets a complicated balance of factors such as absorption, excretion and detoxification (Abdel-Aal, 1996). On the other hand, this shortened longevity of *P. gossypiella* adult females may be attributed to the effect of the tested CSIs on a hormonal activity because there is a close relation between certain hormones and adult longevity, such as representatives of peptide hormone, lipophilic hormones and bioactive amines as reported for *Drosophila* (Simon *et al.*, 2003; Broughton *et al.*, 2005; Carbone *et al.*, 2006). At least one of the *Drosophila* insulin-linked peptides expressed in the median neurosecretory cells (which produce PTTH) is likely to contribute to the endocrine regulation of longevity (Toivonen and Partridge, 2009). However, the exact mode of action of the tested CSIs on the biochemical sites in adults of *P. gossypiella* is unknown until now!

Pre-oviposition period: In most insects, the pre-oviposition period can be called 'ovarian maturation period' and it may be an informative indicator for the ovarian maturation rate, i.e., the shorter period indicates faster rate and *vice versa*. In the present study, treatment of 1-day old eggs of *P. gossypiella* with Noviflumuron led to a slight prolongation or shortening of pre-oviposition period, depending on the concentration while treatment of eggs with Novaluron led to a slight prolongation of such period, in a dose-dependent manner. The present result of prolongation corroborated with those reported results of prolonged period after treatment of newly hatched larvae of the same lepidopterous insect with Diflubenzuron, Hexaflumuron and Chlorfluazuron (Kandil *et al.*, 2005, Kandil *et al.*, 2013), LC₅₀ values of Chromafenozide and Diflubenzuron (Salem, 2015), LC₅₀ of Teflubenzuron (El-Khayat *et al.*, 2015) and after treatment of newly hatched or full grown larvae with different concentrations of Novaluron (Hassan *et al.*, 2017) or after treatment of 1- and 2-day old eggs with LC₅₀ of Hexaflumuron (El-Barkey *et al.*, 2009). It was in agreement, also, with those reported prolongation of the pre-oviposition period in other insects, such as *S. littoralis*, after larval treatment with Diflubenzuron (Aref *et al.*, 2010) and *Ephesia kuehniella*, after larval treatment with Tebufenozide (Bouzera and Soltani-Mazouni, 2014).

On the other hand, the shortened pre-oviposition period after egg treatment with certain concentrations of Noviflumuron, in the present study on *P. gossypiella*, agreed with those reported results of shortened period in the same insect, after treatment of 1- and 2-day old eggs with LC₅₀ of Radiant (Spintoram, bacteria-based product) (El-Barkey *et al.*, 2009) and after treatment of newly hatched larvae with Diflubenzuron (Rashad *et al.*, 2006) as well as after larval

treatment of *D. antique* with a dose of 100 mg kg⁻¹ of Pyriproxyfen (Zhou *et al.*, 2016).

Many lepidopterous species have a relatively short, non-feeding adult stage, which requires the adult female to emerge with most of her eggs ready to be fertilized and oviposited within hours. This life style constrains these insects to a program of ovarian organogenesis and follicle development that must occur at stages earlier than in other insects. The determinants required for germ cell formation are similar in moths, but there are spatial differences in their localization within the presumptive germ band (Richard *et al.*, 1998). In the light of this information, retarding effect of Noviflumuron (at certain concentrations) and Novaluron on the ovarian maturation in *P. gossypiella*, in the present study, may be understood by influenced germ band or the number of germ cells formed in the embryo (Hodin and Riddiford, 1998). However, the exact mode of retarding action is unfortunately available right now but interference of the tested CSIs with the hormonal regulation needs further investigation in the foreseeable future.

Oviposition period: In respect of another important compartment of adult longevity, oviposition period (reproductive life-time), scarcely reported results have been seen in the available literature. According to the few reported results, oviposition period in the adult females of *P. gossypiella* had been shortened after treatment of newly hatched larvae with Chlorfluazuron (Kandil *et al.*, 2005), Diflubenzuron (Rashad *et al.*, 2006), Hexaflumuron and Chlorfluazuron (Kandil *et al.*, 2013), LC₅₀ of Methomyl (El-Khayat *et al.*, 2015) and after treatment of newly hatched or full grown larvae with Novaluron (Hassan *et al.*, 2017) as well as after treatment of 1- and 2-day old eggs with Radiant (El-Barkey *et al.*, 2009). Result of the present study on *P. gossypiella*, was, to a great extent, concomitant to those reported results, since treatment of 1-day old eggs with sublethal concentrations of Noviflumuron and Novaluron resulted in considerably shortened oviposition period (reproductive life-time). On the contrary, this result disagreed with the reported considerable prolongation of oviposition period in the same lepidopterous insect, after treatment of newly hatched larvae with LC₅₀ of Chromafenozide or Diflubenzuron (Salem, 2015) and Teflubenzuron (El-Khayat *et al.*, 2015) and after treatment of 1- and 2-day old eggs with LC₅₀ of Hexaflumuron (El-Barkey *et al.*, 2009).

On the basis of shortened oviposition period of *P. gossypiella* after treatment of 1-day old eggs with Noviflumuron and Novaluron, in the current investigation, these CSIs exerted tremendously enforcing actions on the ovipositing females to lay eggs quickly. This accelerated oviposition may be attributed to a physiological behaviour of the ovipositing adult females to avoid a long time interval under stress of the tested CSIs as xenobiotic factors.

Post- oviposition period: Depending on the currently available literature, very scarce studies have examined the effects of IGRs on post-oviposition period, the last compartment of adult life in insects. With regard to the present experimental insect, *P. gossypiella*, the post-oviposition period was significantly prolonged after treatment of larvae with Hexaflumuron and Chlorfluazuron (Kandil *et*

al., 2013) or treatment of 1- and 2-day old eggs with LC₅₀ of Hexaflumuron (El-Barkey *et al.*, 2009). Results of the present study were, to some extent, in agreement with those reported results, since treatment of 1-day old eggs of *P. gossypiella* with Noviflumuron resulted in pronouncedly prolonged post-oviposition period but slightly prolonged period was recorded after treatment of similar eggs with by Novaluron. In contrast, our result disagreed with those reported results of shortened post-oviposition period of the same insect after treatment of 1- and 2-day old eggs with LC₅₀ of Radiant (El-Barkey *et al.*, 2009) and diverse effect of Novaluron, after treatment of larvae (Hassan *et al.*, 2017). Unfortunately, there is no acceptable interpretation for this effect right now!!

4.2. Disrupted Reproductive Potential of *P. gossypiella* by CSIs

Reproduction in insects is mainly controlled by corpus allatum hormone (juvenile hormone, JH), which is also responsible for protein metabolism, and is specifically needed for egg maturation. The insect growth regulators (IGRs) have been found to render treated insects either sterile or less fecund (Ghoneim *et al.*, 2014). The IGR-treated insects may develop as morphologically deformed adults who would be non-viable or at least their reproductive capacity is reduced (Williams and Amos 1974). However, effects of IGRs on the insect reproduction can be grouped into the following categories: i) reproductive behaviour, ii) oviposition, iii) hatchability of eggs (ovicidal and embryocidal), and iv) sterilization of adults (Mondal and Parween, 2000). On the other hand, ecdysteroids have essential functions in controlling the processes involved in insect reproduction, i.e., vitellogenesis, ovulation of matured eggs and spermatocyte growth (Wigglesworth, 1984; Hagedorn, 1985). In the present study on *P. gossypiella*, treatment of 1-day old eggs with the highest concentration of Noviflumuron (4.0 ppm) or Novaluron (1.0 ppm) led to a failure of the adult females, in spite of mating with normal males, to lay eggs. On the other hand, various degrees of perturbed reproductive potential could be recorded at other concentrations of the present CSIs.

4.2.1. Inhibited oviposition efficiency

In insects, the oviposition rate can be used as an informative indicator for the oviposition efficiency. In the present study on *P. gossypiella*, treatment of 1-day old eggs with sublethal concentrations of Noviflumuron or Novaluron resulted in drastically prohibited oviposition efficiency, since the oviposition rate was severely regressed, in a dose-dependent course. This result was coincided with the reported inhibition of oviposition efficiency of the same lepidopterous insect after treatment of newly hatched and full grown larvae with Novaluron (Hassan *et al.*, 2017) as well as reported results for other insects, such as *S. littoralis* by Tebufenozide (Bakr *et al.*, 2005), Flufenoxuron (Bakr *et al.*, 2010) and Novaluron (Ghoneim *et al.*, 2014); *Schistocerca gregaria* by Flufenoxuron and lufenuron (Soltani-Mazouni and Soltani, 1994) or Tebufenozide (Al-Dali *et al.*, 2008); *Plodia interpunctella* by the ecdysteroid agonist RH-5849 (Smagghe and Degheele, 1994) and *Callosobruchas maculatus* by Cyromazine (Al-Mekhlafi *et al.*, 2011). In contrast, the present result disagreed with the stimulated oviposition of *Gryllus bimaculatus* by some ecdysteroid agonists (Behrens and Hoffmann, 1983).

The prohibited oviposition efficiency, in the current study, may be explained as a result of inhibition of ovarian DNA synthesis or the interference of Noviflumuron and Novaluron

with vitellogenesis in *P. gossypiella* via certain biochemical processes, as will be mentioned later. However, these CSIs may exert a reverse action to those exerted by the ecdysteroid agonists which stimulate the neurosecretory cells to release a myotropic ovulation hormone (Parween *et al.*, 2001).

Perturbation of the reproductive capacity

The reproductive capacity of an insect can be detected by two major parameters: fecundity (mean number of eggs/female) and fertility (egg hatching % or egg viability).

Prohibited fecundity: The available literature contains many results of prohibited fecundity of several insects after treatment of larvae with various IGRs (and CSIs), such as *S. littoralis* after treatment with Diflubenzuron (Aref *et al.*, 2010), Lufenuron (Gaaboub *et al.*, 2012), Methoxyfenozide (Pineda *et al.*, 2009) and Novaluron (Ghoneim *et al.*, 2014). Also, fecundity of other insect species was reduced by various IGRs, such as *E. kuehniella* by Tebufenozide (Khebbab *et al.*, 2008); *Choristoneura rosaceana* (Sun *et al.*, 2000), *Lobesia botrana* (Saenz-de-Cabezón *et al.*, 2005) and *S. litura* (Shahout *et al.*, 2011) by the ecdysteroid agonist methoxyfenozide; *Leptinotarsa decemlineata* (Farinos *et al.*, 1999) and *Tenebrio molitor* (Taibi *et al.*, 2003) by the ecdysteroid agonist Halofenozide (RH-0345); *S. litura* by Chlorfluazuron (Perveen and Miyata, 2000), *M. domestica* by Lufenuron (Hamadah 2003), *D. koenigi* by Flufenoxuron (Khan and Qamar, 2011); *A. kuehniella* by Diflubenzuron and Hexaflumuron (Ashouri *et al.*, 2014); *P. xylostella* by Pyriproxyfen (Mahmoudvand *et al.*, 2015); *Callosobruchus chinensis* by terpene compounds (α -pinene and β -caryophyllene) (Chaubey, 2015); *T. castaneum* (Gado *et al.*, 2015) and *D. antique* (Zhou *et al.*, 2016) by Lufenuron and *C. cephalonica* by Fenoxycarb (Begum and Qamar, 2016); *etc.*

The present results on *P. gossypiella* corroborated, to a great extent, with the previously reported results, since treatment of 1-day old eggs with sublethal concentrations of Noviflumuron or Novaluron resulted in dramatically reduced fecundity of the successfully reproducing females, in a dose-dependent fashion. Comparatively, Noviflumuron exhibited a stronger reducing effect than Novaluron. These results were, also, in agreement with some of the reported results of dramatically reduced fecundity of the same lepidopterous insect after treatment of 1-day old eggs with Hexaflumuron (El-Barkey *et al.*, 2009) and LC₅₀ values of Lufenuron, Chlorfluazuron and Chromafenozide (Kandil *et al.*, 2012). Also, fecundity of *P. gossypiella* was drastically reduced after treatment of newly hatched larvae with Tebufenozide (El-Khayat *et al.*, 2015), Diflubenzuron (Rashad *et al.*, 2006), Chlorfluazuron (Kandil *et al.*, 2005), Buprofezin (Al-Kazafy, 2013), Hexaflumuron and Chlorfluazuron (Kandil *et al.*, 2013), Chromafenozide (Salem, 2015), as well as Pyriproxyfen, Methoxyfenozide and Lufenuron (Sabry and Abdou, 2016) and Novaluron (Hassan *et al.*, 2017).

On the contrary, the recorded results in the current investigation disagree with some reported results of failure of some IGRs to affect the fecundity of various insects, such as Fenoxycarb against *Apis mellifera* (Thompson *et al.*, 2005), Methoxyfenozide against *S. exigua* (Christian-Lius and Pineda, 2010) and Novaluron and Diflubenzuron against *Halyomorpha halys* (Kamminga *et al.*, 2012). Moreover,

feeding of larvae on leaves treated with Methoxyfenozide enhanced the fecundity of *S. littoralis* (Ishaaya *et al.*, 1995). However, these diverse effects can be attributed to the different modes of action of IGRs, different susceptibilities of the insect species, time of treatment and other factors.

The drastically prohibited fecundity of *P. gossypiella*, after treatment of 1-day old eggs with Noviflumuron and Novaluron, in the present study, may be due to its interference with one or more processes, from the ovarian follicle development to egg maturation. In some detail, this can be explained by some reasons. First, the tested CSIs may cause some disorders in the ovaries, including cell death in the germarium, resorption of oocytes in the pre-vitellarium and vitellarium, formation of vitellin envelopes and undue proliferation of follicle cells sometimes resulting in malformation of the whole ovary (Lucantoni *et al.*, 2006; Khan *et al.*, 2007). Second, these CSIs may inhibit the development of some ovarioles and/or synthesis and metabolism of proteinaceous constituents during the oogenesis (Salem *et al.*, 1997). Third, the tested CSIs exerted an inhibitory action on the ecdysone activity, threshold of which is essential for the normal oogenesis (Terashima *et al.*, 2005). Fourth, on the basis of hormonal regulation of insect reproduction, the present CSIs may disturb the production and/or function of the gonadotropic hormone (juvenile hormone, JH) responsible for the synthesis of vitellogenins (yolk precursors) and vitellogenesis (Di Ilio *et al.*, 1999). Fifth, eggs may develop normally in ovaries, but they could not be laid, owing to the adversely deformed ovipositor of adult females or to the reduced mechanical strength (Moreno *et al.*, 1994) or their reabsorption before oviposition (Zhou *et al.*, 2016). Sixth, it may be acceptable to suggest that the prohibited fecundity of *P. gossypiella*, in the current work, may be due to inhibitory effects of the tested CSIs on synthesis of both DNA and RNA, suboptimal nutrition owing to reduced feeding, altered mating behaviour as a result of sublethal intoxication, or a combination of factors.

Reduced fertility: Another parameter of the reproductive capacity in insects is fertility (egg viability). In the present study on *P. gossypiella*, treatment of 1-day old eggs with sublethal concentrations of Noviflumuron or Novaluron led to dangerously reduced fertility of the eggs laid by reproducing adult females. Comparatively, Noviflumuron exhibited a stronger reducing effect than Novaluron. This result was in accordance with those reported results of considerably reduced fertility in the same lepidopterous insect after treatment of 1-day old eggs with Hexaflumuron (El-Barkey *et al.*, 2009), Lufenuron, Chlorfluazuron and Chromafenozide (Kandil *et al.*, 2012), LC₅₀ of Chromafenozide and Diflubenzuron (Salem, 2015), Chlorfluazuron and Hexaflumuron (Kandil *et al.*, 2013) or after treatment of newly hatched larvae with Novaluron (Hassan *et al.*, 2017). Also, our result was, to some extent, in agreement with those reported results of reduced fertility of other insects after treatment of larvae with various IGRs, such as *S. littoralis* by Chlorfluazuron (Sammour *et al.*, 2008), Methoxyfenozide (Pineda *et al.*, 2009), Diflubenzuron (Aref *et al.*, 2010), Lufenuron (Gaaboub *et al.*, 2012), Triflumuron (El-Naggar, 2013) and Novaluron (Ghoneim *et al.*, 2014); *S. litura* by Diufenolan (Perveen and Miyata, 2000) and Chromafenozide (Shahout *et al.*, 2011); *T. molitor* by Halofenozide (Taibi *et al.*

al., 2003); *M. domestica* by Diofenolan (Hamadah 2003), *T. castaneum* by Novaluron (Kostyukovsky and Trostanelsky, 2004); *E. kuehniella* by Tebufenozide (Khebbab *et al.*, 2008); *D. koenigi* by Flufenoxuron (Khan and Qamar, 2011), *C. maculata* by Cyromazine (Al-Mekhlafi *et al.*, 2011), *A. kuehniella* by Diflubenzuron and Hexaflumuron (Ashouri *et al.*, 2014); *etc.*

For explicating the fertility reduction in *P. gossypiella* by Noviflumuron and Novaluron, in the present study, some suggestions can be provided herein. First, maturation of the insect eggs depends basically on the vitellogenins, precursor materials of vitellins including proteins, lipids and carbohydrates, all of which are necessarily required for the embryonic development (Soltani and Mazouni, 1992; Chapman, 1998). These materials are synthesized primarily by fat body during the immature stages (Telfer, 2009) or by the ovary *in situ* (Indrasith *et al.*, 1988). Wherever the site of vitellogenin synthesis, the tested CSIs may disturb their production and/or accumulation in adult females of *P. gossypiella* leading to reduction of fertility. Second, the tested compounds may indirectly affect the fertility *via* the disruption of opening of the intracellular spaces in follicular epithelium or generally inhibition the role of JH (gonadotropic hormone) responsible for the regulation of vitellogenesis (Davey and Gordon, 1996). Third, the fertility reduction may be due to the penetration of residual amounts of the present CSIs into *P. gossypiella* eggs and disturbance of embryonic cuticle synthesis. So, the fully mature embryos had weakened chitinous mouth parts that were insufficiently rigid to perforate the surrounding vitellin membrane for hatching process (Sallam, 1999; Sammour *et al.*, 2008). Fourth, fertility reduction in *P. gossypiella*, in the current study, may be due to serious effects of the present CSIs on the survival of developing embryos at certain stages. Fifth, because the molecular studies revealed the effects of some IGRs on insect reproduction owing to the perturbation of gene expression in the hierarchy cascade of vitellogenesis and/or choriogenesis (Sun *et al.*, 2003), the tested CSIs may interfere with the gene expression resulting in a reduction of the developed embryos in *P. gossypiella*, in the present study.

Retarded embryonic development

In insects, incubation period can be used as a valuable indicator of the embryonic developmental rate, i.e., longer period usually denotes slower rate and *vice versa*. In the present study on *P. gossypiella*, treatment of 1-day old eggs with sublethal concentrations of Noviflumuron or Novaluron led to considerably prolonged incubation period of eggs laid by the successfully reproducing females, denoting severely retarding effect of both CSIs on the embryonic developmental rate. The present result corroborated with the scarcely reported results, in the available literature, concerning a similar retarding action of some IGRs on the embryonic development of the same lepidopterous insect, after treatment of 0-3-day old eggs with LC₅₀ of Hexaflumuron (El-Barkey *et al.*, 2009) and after treatment of larvae with LC₅₀ values of Lufenuron, Chlorfluazuron or Chromafenozide (Kandil *et al.*, 2012) and Novaluron (Hassan *et al.*, 2017). Also, such period was prolonged after treatment of larvae of *C. maculata* with Cyromazine (Al-Mekhlafi *et al.*, 2011) and *S. littoralis* with Novaluron (Ghoneim *et al.*, 2014) as well as after treatment

of 1- and 3-day old egg masses of the latter insect with LC₅₀ of sesame oil (Khedr, 2016). The delayed embryonic development in *P. gossypiella* after treatment of 1-day old eggs with Noviflumuron and Novaluron, in the present study, may be due to their effects on ecdysteroids responsible for the regulation of embryogenesis at certain stages, especially those originating from the ovary *in situ* (Chapman, 1998).

5. CONCLUSIONS

In the light of the recoded results, Noviflumuron and Novaluron exhibited long-term disruptive effects on the adult performance and reproductive capacity of the pink bollworm *P. gossypiella* leading to a considerable reduction of its population. Therefore, the tested compounds, especially Noviflumuron, may be potential IGRs being involved in the integrated control program against this worldwide serious pest.

6. ACKNOWLEDGEMENTS

Authors greatly thank Laboratory of Pesticides, Agricultural Research Center, Doqqi, Giza, Egypt, for providing the Benzoylphenylurea Noviflumuron.

7. REFERENCES

- Abdel-Aal, A.E. 1996. Biological, histological and physiological effects of some insect growth regulators on the greasy cutworm, *Agrotis ipsilon* (Lepidoptera: Noctuidae). M.Sc. Thesis; Faculty of Science, Cairo University Egypt, 183pp.
- Abd-El-Aziz, H.S. and Sayed, S.Z. 2014. Effects of certain insecticides on eggs of *Spodoptera littoralis*. *Egypt. J. Agric. Res.*, **92(3)**: 875-884.
- Abd-Elhady, H.K. and Abdel-Aal, A.A. 2011. Insecticides resistance detection in field-collected populations of *Pectinophora gossypiella* (Saunders). *Am. J. Biochem. Molec. Biol.*, **1(4)**: 337-348.
- Abd El-Hafez, A., Metwally, A.G. and Saleh, M.R.A. 1982. Rearing pink bollworm *Pectinophora gossypiella* (Saund.) on kidney bean diet in Egypt. (Lepidoptera: Gelechiidae). *Res. Bull., Fac. Agric., Zagazig Univ.*, **576**: 10 pp.
- Al-Dali, A.G., Ghoneim, K.S., Bakr, R.F., Bream, A.S. and Tanani, M.A. 2008. Egg productivity of *Schistocerca gregaria* (Orthoptera: Acrididae) as affected by the non-steroidal ecdysone agonist Tebufenozide (RH-5992). *J. Egypt. Acad. Soc. Environ. Develop.*, **9(10)**: 27-38.
- Aliabadi, F.P., Sahragard, A. and Ghadamyari, M. 2016. Lethal and sublethal effects of a chitin synthesis inhibitor, lufenuron, against *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae). *J. Crop Protec.*, **5(2)**: 203-214.
- Al-Mekhlafi, F., Mashaly, A.M., Abdel Mageed, A., Wadaan, M.A. and Al-Mallah, N.M. 2011. Overlap effects of cyromazine concentration, treatment method and rearing temperature on the Southern cowpea weevil (*Callosobruchus maculatus* F.) reared on cowpea. *Afr. J. Microbio. Res.*, **5(32)**: 5848-5853.
- Al-Sharook, Z., Balan, K., Jiang, Y. and Rembold, H. 1991. Insect growth inhibitors from two tropical Meliaceae: Effects of crude seed extracts on mosquito larvae. *J. App. Entomol.*, **111**: 425-430.

- Alyokhin, A., Guillemette, R. and Choban, R. 2009. Stimulatory and suppressive effects of Novaluron on the Colorado potato beetle reproduction. *J. Econ. Entomol.*, **102**(6): 2078-2083.
- Ameen A, Kaakeh W, Wang Ch, and Bennett G. 2002. Laboratory and field efficacy of Noviflumuron formulations against the German cockroach. In Proceedings of 4th International Conference on Urban Pests, (Jones, S.C., Zhai, J. and Robinson, W.H., eds.). *Pocahontas Press, Charleston, SC*, pp.: 147–153.
- Ameen, A., Wang, C., Kaakeh, W., Bennett, G.W., King, J.E. and Karr, L.L. 2005. Residual activity and population effects of noviflumuron for German cockroach (Dictyoptera: Blattellidae) control. *J. Econ. Entomol.*, **98**: 899–905.
- Aref, S.A., Bayoumi, O.Ch. and Soliman, H.A.B. 2010. Effect of certain insecticides on the biotic potential of the cotton leafworm, *Spodoptera littoralis* (Boisd.). *Egypt. J. Agric. Res.*, **88**(1): 31-40.
- Ashouri, S., Pourabad, R.F. and Ebadollahi, A. 2014. The effect of diflubenzuron and hexaflumuron on the last larval instars of the Mediterranean flour moth *Anagasta kuehniella* (Zeller) (Lepidoptera: Pyralidae) under laboratory conditions. *Arch. Phytopathol. Plant Protec.*, **47**(1): 75-81.
- Bakr, R.F.A., Guneidy, N.A.M. and El-Bermawy, S.M. 2005. Toxicological and biological activities of three IGRs against fourth larval instar of the cotton leafworm *Spodoptera littoralis* (Boisd.). *J. Egypt. Acad. Environ. Develop.*, **6**(4): 103-132.
- Bakr, R.F.A., El-barky, N.M., Abd Elaziz, M.F., Awad, M.H. and Abd El-Halim, H.M.E. 2010. Effect of chitin synthesis inhibitors (flufenoxuron) on some biological and biochemical aspects of the cotton leaf worm *Spodoptera littoralis* Bosid. (Lepidoptera: Noctuidae). *Egypt. Acad. J. Biolog. Sci.*, **2**(2): 43-56.
- Barazani, A. 2001. Rimon, an IGR insecticide. *Phytoparasitica*, **29**: 59-60.
- Begum, R. and Qamar, A. 2016. Fenoxycarb- a potent inhibitor of metamorphosis and reproduction in Rice Moth, *Corcyra cephalonica* (Stainton). *J. Entomol. Zool. Studies*, **4**(4): 572-577.
- Behrens, W. and Hoffman, K.H. 1983. Effects of exogenous ecdysteroids on reproduction in cricket *Gryllus bimaculatus*. *Int. Invertr. Reprod. Develop.*, **6**: 149-159.
- Benseba, F., Kilani-Morakchi, S., Aribi, N. and Solatani, N. 2015. Evaluation of pyriproxyfen, a juvenile hormone analog, on *Drosophila melanogaster* (Diptera: Drosophilidae): insecticidal activity, ecdysteroid contents and cuticle formation. *Eur. J. Entomol.*, **112**(4): 625–631. doi: 10.14411/eje.2015.084
- Bouzera, H. and Soltani-Mazouni, N. 2014. Comparative effects of two moulting hormone agonists (Methoxyfenozide and Tebufenozide) on the Mediterranean flour moth *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae): ecdysteroids amounts of testes and reproductive events. *World App. Sci. J.*, **31**(11): 1903-1910.
- Broughton, S.J., Piper, M.D., Ikeya, T., Bass, T.M., Jacobson, J., Driege, Y., Martinez, P., Hafen, E., Withers, D.J., Leever, S.J. and Partridge, L. 2005. Longer lifespan, altered metabolism, and stress resistance in *Drosophila* from ablation of cells making insulin-like ligands. *Proc. Natl. Acad. Sci. U.S.A.*, **102**: 3105–3110.
- Carbone, M.A., Jordan, K.W., Lyman, R.F., Harbison, S.T., Leips, J., Morgan, T.J., DeLuca, M., Awadalla, P. and Mackay, T.F. 2006. Phenotypic variation and natural selection at catsup, a pleiotropic quantitative trait gene in *Drosophila*. *Curr. Biol.*, **16**: 912-919.
- Chapman, R.F. 1998. The insects: structure and function. 4th ed. Cambridge: Cambridge University Press.
- Chaubey, M.K. 2015. Biological activities of terpenes against pulse beetle, *Callosobruchus chinensis* (Coleoptera: Bruchidae). *Entomol. App. Sci. Letters*, **2**(1): 50-61.
- Christian-Luis, R. and Pineda, S. 2010. Toxicity and sublethal effects of methoxyfenozide on *Spodoptera exigua* (Lepidoptera: Noctuidae) *J. Econ. Entomol.*, **103**: 662-667.
- Cutler, G.C., Scott-Dupree, C.D., Tolman, J.H. and Harris, C.R. 2005a. Acute and sublethal toxicity of novaluron, a novel chitin synthesis inhibitor, to *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). *Pest Manage. Sci.*, **61**: 1060-1068.
- Cutler, G.C., Tolman, J.H., Scott-Dupree, C.D. and Harris, C.R. 2005b. Resistance potential of Colorado potato beetle (Coleoptera: Chrysomelidae) to novaluron. *J. Econ. Entomol.*, **98**: 1685-1693.
- Cutler, G.C., Scott-Dupree, C.D., Tolman, J.H. and Harris, C.R. 2007. Field efficacy of novaluron for control of Colorado potato beetle (Coleoptera: Chrysomelidae) on potato. *Crop Protec.*, **26**: 760-767.
- Davey, K.G. and Gordon, D.R.B. 1996. Fenoxycarb and thyroid hormones have JH-like effects on the follicle cells of *Locosta migratoria* in vitro. *Arch. Insect Biochem. Physiol.*, **32**: 613- 626.
- Derbalah, A.S., Khidr, A.A., Moustafa, H.Z. and Taman, A. 2014. Laboratory evaluation of some non-conventional pest control agents against the pink bollworm *Pectinophora gossypiella* (Saunders). *Egypt. J. Biol. Pest Control*, **24**(2): 363-368. <http://www.esbcp.org/index.asp>
- Dhadialla, T.S., Carlson, G.R. and Le, D.P. 1998. New insecticides with ecdysteroidal and juvenile hormone activity. *Annu. Rev. Entomol.*, **43**: 545-569.
- Di Ilio, V., Cristofaro, M., Marchini, D., Nobili, P. and Dallai, R. 1999. Effects of a neem compound on the fecundity and longevity of *Ceratitis capitata* (Diptera: Tephritidae). *J. Econ. Entomol.*, **92**: 76-82.
- El-Aswad, A.F. and Aly, M.I. 2007. Screening of some insecticides against the cotton bollworms, *Pectinophora gossypiella* (Saund.) and *Earias insulana* (Boisd.). *J. Pest Cont. Environ. Sci.*, **15**(2): 63-74.
- El-Barkey, N.M., Amer, A.E. and Kandil, M.A. 2009. Ovicidal activity and biological effect of Radiant and hexaflumuron against eggs of pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). *Egypt. Acad. J. Biol. Sci.*, **2**: 23-36.
- El-Khayat, E.F., Rashad, A.M., Abd-El Zaher, T.R., Shams El-Din, A.M. and Salim, H.S. 2015. Toxicological and biological studies of some pesticidal formulations against *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). *American-Eurasian J. Toxicol. Sci.*, **7**(1): 01-06.
- El-Naggar, J.B.A. 2013. Sublethal effect of certain insecticides on biological and physiological aspects of

- Spodoptera littoralis* (Boisd.). *Nature and Science*, **11(7)**: 19-25.
- El-Sheikh, T.A.A. 2002. Effects of application of selected insect growth regulators and plant extracts on some physiological aspects of the black cutworm, *Agrotis ipsilon* (HUF.). Ph.D. Thesis, Fac. Sci., Ain Shams Univ., Egypt, 211pp.
- Fabrick, J.A. and Tabashnik, B.E. 2012. Similar genetic basis of resistance to *Bt*-toxin Cry1Ac in boll-selected and diet-selected strains of pink bollworm. *Plosone*, **7**: 35658.
- Farinos, G.P., Smagghe, G., Tirry, L. and Castañera, P. 1999. Action and pharmacokinetics of a novel insect growth regulator, halofenozide, in adult beetles of *Aubeonymus mariaefrancisciae* and *Leptinotarsa decemlineata*. *Arch. Insect Biochem. Physiol.*, **41**: 201-213.
- Gaaboub, I., Halawa, S. and Rabiha, A. 2012. Toxicity and biological effects of some insecticides, IGRs and Jojoba oil on cotton leafworm *Spodoptera littoralis* (Boisd.). *J. App. Sci. Res.*, **2**: 131-139.
- Gado, P., Salokhe, S.G. and Deshpande, S.G. 2015. Impact of Lufenuron (5.4% EC) on reproductive end points of *Tribolium castaneum*. *World J. Pharmaceut. Res.*, **4(3)**: 1593-1599.
- Ghoneim, K., Tanani, M., Hamadah, Kh., Basiouny, A. and Waheeb, H. 2014. Inhibited reproductive capacity of Egyptian cotton leaf worm *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae) by the chitin synthesis inhibitor Novaluron. *Egypt. Acad. J. Biolog. Sci.*, **7(2)**: 105-118.
- Ghoneim, K., Tanani, M., Hamadah, Kh., Basiouny, A. and Waheeb, H. 2015. Bioefficacy of Novaluron, a chitin synthesis inhibitor, on survival and development of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). *J. Adv. Zool.*, **1(1)**: 24-35.
- Ghoneim, K., Hassan, H.A., Tanani, M.A. and Bakr, N.A. 2017a. Toxic and disruptive effects of Novaluron, a chitin synthesis inhibitor, on development of the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). *Int. J. Entomol. Res.*, **2(2)**: 36-47.
- Ghoneim, K., Hassan, H.A., Tanani, M.A. and Bakr, N.A. 2017b. Deteriorated larval haemogram in the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) by the chitin synthesis inhibitors, Novaluron and Diufenolan. *Int. J. Modern Res. Rev.*, **5(2)**: 1487-1504.
- Hagedorn, H.H. 1985. The role of ecdysteroids in reproduction. In: "Comprehensive Insect Physiology, Biochemistry and Pharmacology" (Kerkut, G.A. and Gilbert, L.I., eds.). Pergamon, Oxford, vol. **8**, pp.: 205-262.
- Hamadah, Kh.Sh. 2003. Physiological and biochemical effects of IGRs and plant extracts on the house fly *Musca domestica*. M.Sc. Thesis Fac. Sci., Al-Azhar Univ., Cairo, Egypt, 210pp.
- Hamadah, Kh., Tanani, M., Ghoneim, K., Basiouny, A. and Waheeb, H. 2015. Effectiveness of Novaluron, chitin synthesis inhibitor, on the adult performance of Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). *Int. J. Res. Studies Zool.*, **1(2)**: 45-55.
- Hammock, C.D. and Quisted, G.B. 1981. Metabolism and mode of action of juvenile hormone, juvenoids and other insect growth regulators. In "Progress in pesticide Biochemistry" (Hutson, D.H. and Roberts, T.R. eds.), John Wiley & Sons Ltd., **1**: 1-85.
- Hassan, H.A., Ghoneim, K., Tanani, M.A. and Bakr, N.A. 2017. Impairing effectiveness of the chitin synthesis inhibitor, Novaluron, on adult performance and reproductive potential of the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). *J. Entomol. Zool. Studies*, **5(2)**: 581-592.
- Hodin, J. and Riddiford, L.M. 1998. The ecdysone receptor and ultraspiracle regulate the timing and progression of ovarian morphogenesis during *Drosophila metamorphosis*. *Develop. Genes Evol.*, **208**: 304-317.
- Hoffman, K.H. and Lorenz, M.W. 1998. Recent advances in hormones in insect pest control. *Phytoparasitica*, **26(4)**: 323-330.
- Indrasith, L., Sasaki, S.T., Yaginuma, T. and Yamashita, O. 1988. The occurrence of premature form of egg-specific protein in vitellogenic follicles of *Bombyx mori*. *J. Comp. Physiol.*, **158**: 1-7.
- Ishaaya, I. and Horowitz, A.R. 2002. Novaluron (Rimon) a novel IGR: its biological activity and importance in IPM programs. *Phytoparasitica*, **30**: 203.
- Ishaaya, I., Yablonski, S. and Horowitz, A.R. 1995. Comparative toxicity of two ecdysteroid agonists, RH-2485 and RH-5992, on susceptible and pyrethroid-resistant strains of the Egyptian cotton leafworm, *Spodoptera littoralis*. *Phytoparasitica*, **23**: 139-145.
- Ishaaya, I., Kontsedalov, S., Masirov, D. and Horowitz, A.R. 2001. Bio-rational agents-mechanism, selectivity and importance in IPM programs for controlling agricultural pests. *Med. Landbouww Rijksuniv Gent*, **66**: 363-374.
- Ishaaya, I., Horowitz, A.R., Tirry, L. and Barazani, A. 2002. Novaluron (Rimon), a novel IGR: mechanism, selectivity and importance in IPM programs. *Proc. Int. Symp. Crop Protect. Med. Fac. Landbouww Univ., Gent*, **67**: 617-626.
- Ishaaya, I., Kontsedalov, S. and Horowitz, A.R. 2003. Novaluron (Rimon), a novel IGR: Potency and cross-resistance. *Arch. Insect Biochem. Physiol.*, **54**: 157-164.
- Jimenez-Peydro, R., Gimeno-Martos, C., Lopez-Ferrer, J., Serrano-Delgado, C. and Moreno-Mari, J. 1995. Effects of the insect growth regulator, cyromazine, on the fecundity, fertility and offspring development of Mediterranean fruit fly, *Ceratitis capitata* Wied (Diptera, Tephritidae). *J. App. Entomol.*, **119**: 435-438.
- Josephraj Kumar, A., Subrahmanyam, B. and Srinivasan, S. 1999. Plumbagin and azadirachtin deplete haemolymph ecdysteroid levels and alter the activity profiles of two lysosomal enzymes in the fat body of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Eur. J. Entomol.*, **96**: 347-353.
- Kammaing, K.L., Kuhar, T.P., Wimer, A. and Herbert, D.A. 2012. Effects of the insect growth regulators novaluron and diflubenzuron on the brown marmorated stink bug. *Plant Health Progress Online* doi:10.1094/PHP-2012-1212-01-RS.
- Kandil, A.A.M., Abd El-Zhar, T.R. and Rashad, A.M. 2005. Some biological and biochemical effects of chitin synthesis inhibitor on pink bollworm *Pectinophora*

- gossypiella*. *Ann. Agric. Sci. Moshtohor, Egypt*, **43(4)**: 1991-2002.
- Kandil, M.A., Ahmed, A.F. and Moustafa, H.Z. 2012. Toxicological and biochemical studies of lufenuron, chlorfluazuron and chromafenozide against *Pectinophora gossypiella* (Saunders). *Egypt. Acad. J. Biolog. Sci.*, **4(1)**: 37-47.
- Kandil, M.A.A., Salem, M.S. and Adly, A.M. 2013. Biological and biochemical changes in pink bollworm, *Pectinophora gossypiella* after treatment with Hexaflumuron and Chlorfluazuron. *Ann. Agric. Sci. Moshtohor, Egypt*, **51(4)**: 472-437.
- Karr, L.L., Sheets, J.J., King, J.E. and Dripps, J.E. 2004. Laboratory performance and pharmacokinetics of the benzoylphenylurea noviflumuron in eastern subterranean termites (Isoptera: Rhinotermitidae). *J. Econ. Entomol.*, **97**: 593-600.
- Khan, I. and Qamar, A. 2011. Biological activity of andalin (flucycloxuron), a novel chitin synthesis inhibitor, on red cotton stainer *Dysdercus koenigii* (Fabricius). *Frontiers in Life Sciences: Basic and Applied, Biology and Medicine*, **3(2)**: 324-335.
- Khan, I. and Qamar, A. 2012. Andalin, an insect growth regulator, as reproductive inhibitor for the red cotton stainer, *Dysdercus koenigii* (F.) (Hemiptera: Pyrrhocoridae). *Acad. J. Entomol.*, **5(2)**: 113-121.
- Khan, M., Hossain, M.A. and Islam, M.S. 2007. Effects of neem leaf dust and a commercial formulation of a neem compound on the longevity, fecundity and ovarian development of the melon fly, *Bactocera cucurbitae* (Coquillett) and the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). *Pak. J. Biol. Sci.*, **10**: 3656-3661.
- Khan, G.Z., Khan, I., Khan, I.A., Alamzeb Salman, M. and Kalim Ullah 2016. Evaluation of different formulations of IGRs against *Aedes albopictus* and *Culex quinquefasciatus* (Diptera: Culicidae). *Asian Pacific J. Trop. Biomed.*, **6(6)**: 485-491.
- Khatter, H. and Abuldahb, M. 2011. Combined effect of three insect growth regulators on the digestive enzymatic profiles of *Callosobruchus maculatus* (Coleoptera: Bruchidae). *J. Egypt. Soc. Parasitol.*, **41(3)**: 765-76.
- Khebbeb, M.E.H., Gaouaoui, R. and Bendjeddou, F. 2008. Tebufenozide effects on the reproductive potentials of the Mediterranean flour moth, *Ephestia kuehniella*. *Afr. J. Biotech.*, **7(8)**: 1166-1170.
- Khedr, M.A. 2016. Ovicidal activity of *Sesamum indicum* (L.) oil against *Spodoptera littoralis* (Boisd.). *Arch. Phytopathol. Plant Protec.*, **49(5-6)**: 95-110.
- Khidr, A.A., Moawad, G.M., Desuky, W.M.H. and Raslan, S.A. 1996. Effect of some synthetic pyrethroids on bollworm larvae in cotton fields. *Egypt. J. Agric. Res.*, **74(1)**: 123-132.
- Kim, K.S., Chung, B.J. and Kim, H.K. 2000. DBI-3204: A new benzoylphenyl urea insecticide with a particular activity against whitefly. *Brighton Crop Prot. Conf.*, **1**: 41-46.
- King, J.E. 2005. Ovicidal activity of noviflumuron when fed to adult German cockroaches (Dictyoptera: Blattellidae). *J. Econ. Entomol.*, **98**: 930-932.
- Kostyukovsky, M. and Trostanetsky, A. 2004. The effect of a new chitin synthesis inhibitor, Novaluron, on various developmental stages of *Tribolium castaneum* (Herbst). *J. stored Prod. Res.*, **42(2)**: 136-148.
- Kranthi, K.R. 2015. Pink bollworm strikes Bt cotton. *Cotton Statistics News*, **35**: 1-6.
- Liu, Y.D., Wu, K.M. and Guo, Y.Y. 2009. Population structure and introduction history of the pink bollworm, *Pectinophora gossypiella*, in China. *Entomologia Experimentalis et Applicata*, **130**: 160-172.
- Lucantoni, L., Giusti, F., Cristofaro, M., Pasqualini, L., Esposito, F., Lupetti, P. and Habluetzel, A. 2006. Effects of a neem extract on blood feeding, oviposition and oocyte ultrastructure in *Anopheles stephensi* Liston (Diptera: Culicidae). *Tissue Cell*, **38**: 361-371.
- Luna, J.-C., Robinson, V.-A., Martinez, A.-M., Schneider, M.-I., Figueroa, J.-I., Smaghe, G., Vinuela, E., Budia, F. and Pineda, S. 2011. Long-term effects of methoxyfenozide on the adult reproductive processes and longevity of *Spodoptera exigua* (Lepidoptera: Noctuidae). *J. Econ. Entomol.*, **104(4)**: 1229-1235. doi: 10.1603/EC10450.
- Mahmoudvand, M., Abbasipour, H., SheikhiGarjan, A. and Bandani, AR. 2012. Decrease in pupation and adult emergence of *Plutella xylostella* (L.) treated with hexaflumuron. *Chilean J. Agric. Res.*, **72(2)**: 206-211.
- Mahmoudvand, M., Moharramipour, S. and Iranshahi, M. 2015. Effects of pyriproxyfen on life table indices of *Plutella xylostella* in multigenerations. *Psyche*, Article ID **453701**, 7
- Malhat, F.M., Loutfy, N.M. and Ahmed, M.T. 2014. Dissipation kinetics of novaluron in tomato, an arid ecosystem pilot study. *Toxicol. Environ. Chem.*, **96(1)**: 41-47. doi: 10.1080/02772248.2014.911538
- Martins, F. and Silva, I.G. 2004. Avaliação da atividade inibidora do di-flubenzuron na ecdise das larvas de *Aedes aegypti* (Linnaeus, 1762) (Diptera, Culicidae). *Rev. Soc. Bras. Med. Trop.*, **37**: 135-138.
- Mohammed, A. 2013. Targeting the vacuolar ATPase subunits B and C in pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera; Gelechiidae). *Egypt. J. Genet. Cytol.*, **42**: 365-378.
- Mojaver, M. and Bandani, A.R. 2010. Effects of the insect growth regulator pyriproxyfen on immature stages of sunn pest, *Eurygaster integriceps* Puton (Heteroptera: Scutelleridae). *Munis Entomol. Zool.*, **5(1)**: 187-197.
- Mondal, K. and Parween, S. 2000. Insect growth regulators and their potential in the management of stored-product insect pests. *Integr. Pest Manage. Rev.*, **5**: 255-295.
- Monsanto 2010. Cotton in India. http://monsanto.com/news_views/Pages/india-pink-bollworm.aspx (subsequently revised to "Pink bollworm resistance to GM cotton in India").
- Moreno, D.S., Martinez, A.J. and Riviello, M.S. 1994. Cyromazine effects on the reproduction of *Anastrepha ludens* (Diptera: Tephritidae) in the laboratory and in the field. *J. Econ. Entomol.*, **87**: 202-211.
- Moroney, M.J. 1956. Facts from figures (3rd ed.). Penguin Books Ltd., Harmondsworth. Middle Sex.
- Parween, S., Faruki, S.I. and Begum, M. 2001. Impairment of reproduction in the red flour beetle, *Tribolium castaneum* (Herbst.) (Coleoptera: Tenebrionidae) due to

- larval feeding on triflumuron-treated diet. *J. App. Entomol.*, **125**: 1-4.
- Patil, S.B. 2003. Studies on management of cotton pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). Ph.D. Thesis. University of Agricultural Sciences, Dharwad (India).
- Perveen, F. and Miyata, T. 2000. Effects of sublethal dose of Chlorfluazuron on ovarian development and oogenesis in the common cutworm *Spodoptera litura* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.*, **93**(5): 1131-1137.
- Pineda, S., Budia, F., Schneider, M.I., Gobbi, A., Vinuela, E., Valle, J. and Estal, P. del 2004. Effects of two biorational insecticides, spinosad and methoxyfenozide, on *Spodoptera littoralis* (Lepidoptera: Noctuidae) under laboratory conditions. *J. Econ. Entomol.*, **97**: 1906-1911.
- Pineda, S., Martinez, A.M., Figueroa, J.I., Schneider, M.I., Estal, P del., Vinuela, E., Gomez, B., Smagghe, G. and Budia, F. 2009. Influence of azadirachtin and methoxyfenozide on life parameters of *Spodoptera littoralis*. *J. Econ. Entomol.*, **102**(4): 1490-1496.
- Portilla, M., Snodgrass, G. and Luttrell, R. 2012. A novel bioassay using a non-autoclaved solid *Lygus* diet to evaluate the effect of *Beauveria bassiana* and the insect growth regulator novaluron on tarnished plant bug, *Lygus lineolaris*, 3rd international *Lygus* symposium, Scottsdale, Arizona, USA.
- Radwan, E.M.M. and El-Malla, M.A. 2015. Toxicity of insecticides against the laboratory and field moths *Pectinophora gossypiella* (Saunders). *Egypt. Sci. J. Pestic.*, **1**(1): 9pp.
- Ragaei, M. and Sabry, K.H. 2011. Impact of spinosad and buprofezin alone and in combination against the cotton leafworm, *Spodoptera littoralis* under laboratory conditions. *J. Biopestic.*, **4**(2): 156-160.
- Rashad, A.M., Hewady, M.A.A. and Kandil, M.A.A. 2006. Effect of Neemazal, Spinosad and Dimilin on some biological and physiological activities of pink bollworm *Pectinophora gossypiella* (Saund.). *Ann. Agric. Sci., Moshtohor, Egypt*, **44**(1): 304-319.
- Rashad, A.M., El-Khayat, E.F., Abd-El Zaher, T.R., Shams El-Din, A.M. and Salim, H.S. 2015. Biochemical studies of some pesticidal formulations against *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae). *American-Eurasian J. Agric. Environ. Sci.*, **15**(3): 303-307.
- Reinke, M.D. and Barrett, B.A. 2007. Fecundity, fertility and longevity reductions in adult oriental fruit moth (Lepidoptera: Tortricidae) exposed to surfaces treated with the ecdysteroid agonists tebufenozide and methoxyfenozide. *J. Entomol. Sci.*, **42**: 457-466.
- Richard, D.S., Watkins, N.L., Serafin, R.B. and Gilbert, L.I. 1998. Ecdysteroids regulate yolk protein uptake by *Drosophila melanogaster* oocytes. *J. Insect Physiol.*, **44**: 637-644.
- Rose, R.I. 2001. Pesticides and public health: integrated methods of mosquito management. *Emerg. Infect. Dis.*, **7**(1): 17-23.
- Sabry, K.H. and Abdel-Aziz, N.F. 2013. Resistance and enzyme assessment of the pink bollworm, *Pectinophora gossypiella* (Saunders) to Spinosad. *J. Animal Plant Sci.*, **23**(1): 136-142.
- Sabry, K.H. and Abdou, G.Y. 2016. Biochemical and toxic characterization of some insect growth regulators to the pink bollworm, *Pectinophora gossypiella* (Saunders). *American-Eurasian J. Sustainable Agric.*, **10**(1): 8-14.
- Saenz-de-Cabezón, I.F.J., Marco, V., Salmo, F.G. and Perez-Moreno, I. 2005. Effects of methoxyfenozide on *Lobesia botrana* Den and Schiff (Lepidoptera: Tortricidae) egg, larval and adult stages. *Pest Manage. Sci.*, **11**: 1133-1137.
- Salama, M.A., Abd El-Baki, M.A., El-Naggar, J.B.A. and El-Naggar, E.Y. 2013. Efficiency of some insecticides sequence on cotton bollworms and histopathological effects of some biocides on pink bollworm larvae. *Egypt. J. Agric. Res.*, **91**(2): 429-448.
- Salem, M.S.M. 2015. Latent effect of different compounds on *Pectinophora gossypiella* (Saunders). *J. Plant Prot. and Path., Mansoura Univ., Egypt*, **6**(2): 269-279.
- Salem, H., Smagghe, G. and Degheele, D. 1997. Effects of tebufenozide on oocyte growth in *Plodia interpunctella*. *Medical. Faculty. Landbouww. Gent Univ.*, **62**(1): 9-13.
- Sallam, M.H. 1999. Effect of Diflubenzuron on embryonic development of the acridid, *Heteracris littoralis*. *J. Egypt. Ger. Soc. Zool.*, **30**(E): 17-26.
- Sammour, E.A., Kandit, M.A. and Abdel-Aziz, N.F. 2008. The reproductive potential and fate of chlorfluazuron and lufenuron against cotton leafworm, *Spodoptera littoralis* (Boisd). *American-Eurasian J. Agric Environ. Sci.*, **4**(1): 62-67.
- Shahout, H.A., Xu, J.-X., Qiao, J. and Jia, Q.-D. 2011. Sublethal effects of methoxyfenozide, in comparison to chlorfluazuron and beta-cypermethrin, on the reproductive characteristics of common cutworm *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae). *J. Entomol. Res. Soc.*, **13**(3): 53-63.
- Sheets, J. and Karr, L. 2001. Kinetics of uptake, clearance, transfer, and metabolism of noviflumuron in termites (*Reticulitermes flavipes*). The Annual Meeting of the Entomological Society of America, San Diego, California. December 9-12, 2001. (December 28, 2001).
- Simon, A.F., Shih, C., Mack, A. and Benzer, S. 2003. Steroid control of longevity in *Drosophila melanogaster*. *Sci.*, **299**: 1407-1410.
- Singh, A. and Tiwari, S.K. 2016. Role of Fenoxycarb, a juvenile hormone analogue, on the developmental stages of rice-moth, *Corcyra cephalonica* Staint. (Lepidoptera: Pyralidae). *Int. J. Zool. Investig.*, **2**(2): 267-280.
- Smagghe, G. and Degheele, D. 1994. The significance of pharmacokinetics and metabolism to the biological activity of RH-5992 (tebufenozide) in *Spodoptera exempta*, *Spodoptera exigua* and *Leptinotarsa decemlineata*. *Pest. Biochem. Physiol.*, **49**: 224-234.
- Smith, M.S., Karr, L.L., King, J.E., Kline, W.N., Sbragia, R.J., Sheets, J.J. and Tolley, M. 2002. Noviflumuron activity in household and structural insect pests. In: Proceedings of 4th International Conference on Urban Pests (Jones, S.C., Zhai, J. and Robinson, W.H., eds.). Pocahontas Press, Charleston, SC, pp.: 345-354.
- Soltani, N. and Mazouni, A. 1992. Diflubenzuron and oogenesis in the codling moth, *Cydia pomonella*. *Pesti. Sci.*, **34**: 257-261.

- Soltani-Mazouni, N. and Soltani, N. 1994. Diflubenzuron affected DNA synthesis in the ovaries of *Tenebrio molitor*. *J. Invert. Reprod. Devel.*, **25**: 19-21.
- Soltani, N., Besson, M.T. and Delachambre, J. 1984. Effects of diflubenzuron on the pupal – adult development of *Tenebrio molitor* (L.) (Coleoptera: Tenebrionidae): growth and development cuticle secretion, epidermal cell density, and DNA synthesis. *Pestic. Biochem. Physiol.*, **21**:256-264.
- Sun, X., Barrett, B.A. and Biddinger, D.J. 2000. Fecundity and fertility reductions in adult leafrollers exposed to surfaces treated with the ecdysteroid agonists tebufenozide and methoxyfenoside. *Entomol. Exp. App.*, **94**: 75-83.
- Sun, X., Song, Q. and Barrett, B. 2003. Effect of ecdysone agonists on vitellogenesis and the expression of EcR and USP in codling moth (*Cydia pomonella*). *Arch. Insect Biochem. Physiol.*, **52**:115-129.
- Taibi, F., Smagghe, G., Amrani, L. and Soltani-Mazouni, N. 2003. Effect of ecdysone agonist RH-0345 on reproduction of mealworm, *Tenebrio molitor*. *Comp. Biochem. Physiol.*, **135**: 257-267.
- Taleh, M., Pourabad, R.F., Geranmaye, J. and Ebadollahi, A. 2015. Toxicity of Hexaflumuron as an insect growth regulator (IGR) against *Helicoverpa armigera* Hubner (Lepidoptera: Noctuidae). *J. Entomol. Zool. Studies*, **3**(2): 274-277.
- Tanani, A.M. 2001. Study on the effects of certain IGRs and plant extracts on some physiological aspects of the red palm weevil *Rhynchophorus ferrugineus* (Curculionidae: Coleoptera). M.Sc. Thesis, Fac. Sci., Al-Azhar Univ., Egypt, 223pp.
- Tanani, M., Hamadah, Kh., Ghoneim, K., Basiouny, A. and Waheeb, H. 2015. Toxicity and bioefficacy of Cyromazine on growth and development of the cotton leafworm *Spodoptera littoralis* (Lepidoptera: Noctuidae). *Int. J. Res. Studies Zool.*, **1**(3):1-15.
- Tanani, M.A., Ghoneim, K., Hassan, H.A. and Bakr, N.A. 2017. Perturbation of main body metabolites in the pink bollworm *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) by the chitin synthesis inhibitors Novaluron and Diufenolan. *BioBulletin*, **3**(3): (In press).
- Telfer, W.H. 2009. Egg formation in Lepidoptera. *J. Insect Sci.*, **9**: 50. (insectscience.org/9.50).
- Terashima, J., Takaki, K., Sakurai, S. and Bownes, M. 2005. Nutritional status affects 20-hydroxyecdysone concentration and progression of oogenesis in *Drosophila melanogaster*. *J. Endocrinol.*, **187**: 69-79.
- Thompson, H.M., Wilkins, S., Battersby, A.H., Waite, R.J. and Wilkinson, D. 2005. The effects of four insect growth-regulating (IGR) insecticides on honeybee (*Apis mellifera* L.) colony development, queen rearing and drone sperm production. *Ecotoxicol.*, **14**: 757-769.
- Toivonen, J.M. and Partridge, L. 2009. Endocrine regulation of aging and reproduction in *Drosophila*. *Molec. Cell. Endocrinol.*, **299**: 39-50.
- Topozada, A., Abd-allah, S. and El-Defrawi, M.E. 1966. Chemosterilization of larvae and adults of the Egyptian cotton leafworm, *Prodenia litura* by apholate, metepa and hempa. *J. Econ. Entomol.*, **59**: 1125-1128.
- Tunaz, H. and Uygun, N. 2004. Insect growth regulators for insect pest control. *Turkish J. Agric. Forestry*, **28**: 337-387.
- Wang, Ch. and Bennett, W.G. 2006. Efficacy of noviflumuron gel bait for control of the German cockroach, *Blattella germanica* (Dictyoptera: Blattellidae)-laboratory studies. *Pest Manage. Sci.*, **62**:434-439.
- Wang, Q.L. and Liu, T.-X. 2016. Effects of three insect growth regulators on *Encarsia formosa* (Hymenoptera: Aphelinidae), an endoparasitoid of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *J. Econ. Entomol.*, **0**(0): 1-8. doi: 10.1093/jee/tow216
- Wigglesworth, V.B. 1984. *Insect Physiology*. 8th ed., Chapman & Hall, London.
- Williams, D. and Amos, T.G. 1974. Some effects of synthetic juvenile hormones and hormone analogues on *Tribolium castaneum* (Herbs.). *Aust. J. Zool.*, **22**:147-53.
- Zhou, F., Zhu, G., Zhao, H., Wang, Z., Xue, M., Li, X., Xu, H., Ma, X. and Liu, Y. 2016. Sterilization effects of adult-targeted baits containing insect growth regulators on *Delia antiqua*. *Sci. Rep.*, **6**: 32855: 9pp. doi: 10.1038/srep32855.
